

Hygenix, Inc
49 Woodside St.
Stamford, Connecticut 06902
Project: **349 PEPPER RIDGE RD, STAMFORD. CT / NEWFIELD SCHOOL**
Condition of Sample(s) Upon Receipt: Acceptable

Date Collected: 10/29/2018
Date Received: 10/30/2018
Date Analyzed: 10/30/2018
Date Reported: 10/31/2018
Project ID: 18041720
Page 1 of 3

1054 Spore Trap Analysis: SOP 3.8

Client Sample Number	102918-02				102918-01			
Sample Location	MAIN OFFICE				OUTDOOR			
Sample Volume (L)	75				75			
Lab Sample Number	18041720-002				18041720-001			
Spore Identification	Raw Ct	spr/m ³	% Ttl	In/Out	Raw Ct	spr/m ³	% Ttl	In/Out
ascospores	16	853	26	1/1	20	1067	6	-
basidiospores	36	1920	58	1/9	318	16960	93	-
Cladosporium	3	160	5	2/1	2	107	1	-
hyphal elements	-	-	-	-	1	53	<1	-
Penicillium/Aspergillus group	3	160	5	-	-	-	-	-
Smuts,Periconia,Myxomycetes	4	213	6	-	-	-	-	-
	Debris Rating 3				Debris Rating 3			
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m ³				Analytical Sensitivity: 13 spr/m ³			
Comments								
Total *See Footnotes	62	3307	~100%	1/6	341	18187	~100%	-

Client Sample Number	102918-03				102918-01			
Sample Location	ROOM 207				OUTDOOR			
Sample Volume (L)	75				75			
Lab Sample Number	18041720-003				18041720-001			
Spore Identification	Raw Ct	spr/m ³	% Ttl	In/Out	Raw Ct	spr/m ³	% Ttl	In/Out
ascospores	2	107	4	1/10	20	1067	6	-
basidiospores	31	1653	69	1/10	318	16960	93	-
Cladosporium	1	53	2	1/2	2	107	1	-
hyphal elements	1	53	2	1/1	1	53	<1	-
Penicillium/Aspergillus group	6	320	13	-	-	-	-	-
Smuts,Periconia,Myxomycetes	4	213	9	-	-	-	-	-
	Debris Rating 3				Debris Rating 3			
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m ³				Analytical Sensitivity: 13 spr/m ³			
Comments								
Total *See Footnotes	45	2400	~100%	1/8	341	18187	~100%	-

Hygenix, Inc
 49 Woodside St.
 Stamford , Connecticut 06902
 Project: **349 PEPPER RIDGE RD, STAMFORD. CT / NEWFIELD SCHOOL**
 Condition of Sample(s) Upon Receipt: Acceptable

Date Collected: 10/29/2018
 Date Received: 10/30/2018
 Date Analyzed: 10/30/2018
 Date Reported: 10/31/2018
 Project ID: 18041720
 Page 2 of 3

Client Sample Number	102918-04				102918-01			
Sample Location	AUDITORIUM				OUTDOOR			
Sample Volume (L)	75				75			
Lab Sample Number	18041720-004				18041720-001			
Spore Identification	Raw Ct	spr/m ³	% Ttl	In/Out	Raw Ct	spr/m ³	% Ttl	In/Out
ascospores	10	533	9	1/2	20	1067	6	-
basidiospores	86	4587	81	1/4	318	16960	93	-
Cladosporium	5	267	5	3/1	2	107	1	-
hyphal elements	1	53	1	1/1	1	53	<1	-
Penicillium/Aspergillus group	1	53	1	-	-	-	-	-
Smuts,Periconia,Myxomycetes	3	160	3	-	-	-	-	-
	Debris Rating 3				Debris Rating 3			
Analytical Sensitivity	Analytical Sensitivity: 13 spr/m³				Analytical Sensitivity: 13 spr/m³			
Comments								
Total *See Footnotes	106	5653	~100%	1/3	341	18187	~100%	-

Hygenix, Inc
49 Woodside St.
Stamford, Connecticut 06902
Project: **349 PEPPER RIDGE RD, STAMFORD. CT / NEWFIELD SCHOOL**
Condition of Sample(s) Upon Receipt: Acceptable

Date Collected: 10/29/2018
Date Received: 10/30/2018
Date Analyzed: 10/30/2018
Date Reported: 10/31/2018
Project ID: 18041720
Page 3 of 3

Footnotes and Additional Report Information

Debris Rating Table

1	Minimal (<5%) particulate present	Reported values are minimally affected by particulate load.
2	5% to 25% of the trace occluded with particulate	Negative bias is expected. The degree of bias increases directly with the percent of the trace that is occluded.
3	26% to 75% of the trace occluded with particulate	Negative bias is expected. The degree of bias increases directly with the percent of the trace that is occluded.
4	75% to 90% of the trace occluded with particulate	Negative bias is expected. The degree of bias increases directly with the percent of the trace that is occluded.
5	Greater than 90% of the trace occluded with particulate	Quantification not possible due to large negative bias. A new sample should be collected at a shorter time interval or other measures taken to reduce particulate load.

1. Penicillium/Aspergillus group spores are characterized by their small size, round to ovoid shape, being unicellular, and usually colorless to lightly pigmented. There are numerous genera of fungi whose spore morphology is similar to that of the Penicillium/Aspergillus type. Two common examples would be Paecilomyces and Acremonium. Although the majority of spores placed in this group are Penicillium, Aspergillus, or a combination of both. Keep in mind that these are not the only two possibilities.

2. Ascospores are sexually produced fungal spores formed within an ascus. An ascus is a sac-like structure designed to discharge the ascospores into the environment, e.g. Ascobolus.

3. Basidiospores are typically blown indoors from outdoors and rarely have an indoor source. However, in certain situations a high basidiospore count indoors may be indicative of a wood decay problem or wet soil.

4. The colorless group contains colorless spores which were unidentifiable to a specific genus. Examples of this group include Acremonium, Aphanocladium, Beauveria, Chrysosporium, Engyodontium microconidia, yeast, some arthrospores, as well as many others.

5. Hyphae are the vegetative mode of fungi. Hyphal elements are fragments of individual Hyphae. They can break apart and become airborne much like spores and are potentially allergenic. A mass of hyphal elements is termed the mycelium. Hyphae in high concentration may be indicative of colonization.

6. Dash (-) in this report, under raw count column means 'not detected (ND)'; otherwise 'not applicable' (NA).

7. The positive-hole correction factor is a statistical tool which calculates a probable count from the raw count, taking into consideration that multiple particles can impact on the same hole; for this reason the sum of the calculated counts may be less than the positive hole corrected total.

8. Due to rounding totals may not equal 100%.

9. Analytical Sensitivity for each spores is different for Non-viable sample when the spores are read at different percentage. Analytical Sensitivity is calculated as spr/m^3 divided by raw count. $\text{spr}/\text{m}^3 = \text{raw counts} \times (100/\% \text{ read}) \times (1000/\text{Sample volume})$. If Analytical Sensitivity is 13 spr/m^3 at 100% read, Analytical Sensitivity at 50% read would be 27 spr/m^3 , which is 2 times higher. Analytical Sensitivity provided on the report is based on an assumed 100% of the trace being analyzed.

10. Minimum Reporting Limits (MRL) for BULKS, DUSTS, SWABS, and WATER samples are a calculation based on the sample size and the dilution plate on which the organism was counted. Results are a compilation of counts taken from multiple dilutions and multiple medias. This means that every genus of fungi or bacteria recovered can be counted on the plate on which it is best represented.

11. If the final quantitative result is corrected for contamination based on the blank, the blank correction is stated in the sample comments section of the report.


12. The results in this report are related to this project and these samples only.

13. For samples with an air volume of < 100L, the number of significant figures in the result should be considered (2) two. For samples with air volumes between 100-999L, the number of significant figures in the result should be considered (3) three. For example, a sample with a result of 55,443 spr/m^3 from a 75L sample using significant figures should be considered 55,000. The same result of 55,443 from a 150L sample using significant figures should be considered 55,400 spr/m^3 .

14. If the In/Out ratio is greater than 100 times it is indicated >100/1, rather than showing the real value.

Terminology Used in Direct Exam Reporting

Conidiophores are a type of modified hyphae from which spores are born. When seen on a surface sample in moderate to numerous concentrations they may be indicative of fungal growth.



Suzanne S. Blevins, B.S., SM (ASCP)
Laboratory Director